

PAY AS LITTLE AS \$25

for your next prescription of Treximet®

Pay as little as \$ **25***
for your next
prescription of

Treximet®
sumatriptan / naproxen sodium

BIN: 600428
PCN: 06780000
GROUP: 06780014
ID: 50828772410

*This offer will reduce your out-of-pocket cost down to as little as \$25. See Eligibility and Restrictions.

TO PATIENT: Present the coupon to the left with your prescription for Treximet® to your pharmacist and receive savings on your copay. This offer may buy your out-of-pocket cost down to \$25. Maximum program benefit will be dependent upon your insurance and the product quantity purchased. You are responsible for any cost remaining after program benefit is applied. See Eligibility and Restrictions.

TO PHARMACIST: Process a Coordination of Benefits (COB) transaction using your customer's prescription insurance for the primary claim and using BIN: 600428 and PCN: 06780000 for the secondary claim. The secondary claim will buy the out-of-pocket cost down to \$25. Maximum program benefit will be dependent upon patient's insurance and the product quantity purchased. Patient is responsible for any cost remaining after program benefit is applied. Present this card back to your customer for possible future uses. This card can be used for patients who have commercial insurance as well as patients who are cash payers.

FOR PROCESSING QUESTIONS, PLEASE CALL 1-855-830-9254.

ELIGIBILITY AND RESTRICTIONS: This savings card will allow most patients to pay as little as \$25 for their Treximet® prescriptions. Patients with high copay plans or prescription plans based on an annual deductible may have additional responsibility. Maximum benefit is dependent upon patient's individual prescription plan and product quantity purchased. There is no redemption limit. Card is valid for insured patients and cash buyers. Offer valid only in the USA. This offer not valid if your prescription is paid/partially paid by Medicaid, Medicare, Federal or State healthcare programs. PERNIX Therapeutics Holdings, Inc. reserves the right to rescind, revoke or amend this offer without notice. Void where prohibited by law.

If your pharmacist is unable to provide the copay savings at the time you fill your prescription, you can still take advantage of this program if eligible.

- Complete this form with your name and address.
- Circle the product name, date, your name, and amount paid on the original pharmacy receipt. Cash register receipt NOT accepted.
- Mail this form, your pharmacy receipt, and a copy of your Treximet® Savings Card to:
Treximet® Savings Program
2250 Perimeter Park Drive, Suite 300
Morrisville, North Carolina 27560
- In 10-14 days, you will receive a check in the mail.

FIRST NAME _____ LAST NAME _____

ADDRESS _____ SUITE OR APT # _____

CITY _____ STATE _____ ZIP _____

SIGNATURE _____

By my signature, I certify that I meet the Eligibility Criteria listed on this certificate.

For more information visit www.treximet.com.
Please see Important Safety Information below.

PATIENT IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TREXIMET?

TREXIMET may increase your chance of a heart attack or stroke that can lead to death. TREXIMET contains 2 medicines: sumatriptan and naproxen sodium (a nonsteroidal anti-inflammatory drug [NSAID]).

- This risk may happen early in treatment and may increase:
 - with increasing doses of NSAIDs
 - with longer use of NSAIDs.

Do not take TREXIMET right before or after a heart surgery called a "coronary artery bypass graft (CABG)."

Avoid taking TREXIMET after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.

Stop taking TREXIMET and get emergency help right away if you have any of the following symptoms of a heart attack or stroke:

- discomfort in the center of your chest that lasts for more than a few minutes, or that goes away and comes back
- slurred speech
- feeling lightheaded
- severe tightness, pain, pressure, or heaviness in your chest, throat, neck, or jaw
- weakness in one part or on one side of your body
- pain or discomfort in your arms, back, neck, jaw, or stomach
- nausea or vomiting
- shortness of breath with or without chest discomfort
- breaking out in a cold sweat

TREXIMET is not for people with risk factors for heart disease unless a heart exam is done and shows no problem. You have a higher risk for heart disease if you:

- have high blood pressure
- are overweight
- smoke
- have diabetes
- have high cholesterol levels
- have a family history of heart disease

TREXIMET can cause ulcers and bleeding in the stomach and intestines at any time during your treatment.

Ulcers and bleeding can happen without warning symptoms and may cause death.

Your chance of getting an ulcer or bleeding increases with:

- past history of stomach ulcers, or stomach or intestinal bleeding with use of NSAIDs
- older age
- the use of medicines called "corticosteroids," "anticoagulants," and antidepressant medicines called "SSRIs" or "SNRIs"
- longer use
- more frequent use
- drinking alcohol
- having poor health
- advanced liver disease
- bleeding problems
- smoking

TREXIMET may cause serious allergic reactions or serious skin reactions that can be life-threatening. Stop taking TREXIMET and get emergency help right away if you develop:

- sudden wheezing
- rash
- problems breathing or swallowing
- blisters or bleeding of your lips, eye lids, mouth, nose, or genitals
- swelling of your lips, tongue, or body
- fainting
- reddening of your skin with blisters or peeling

TREXIMET should only be used exactly as prescribed, at the lowest dose possible for your treatment, and for the shortest time needed.

TREXIMET already contains an NSAID (naproxen). Do not use TREXIMET with other medicines to lessen pain or fever or with other medicines for colds or sleeping problems without talking to your healthcare provider first, because they may contain an NSAID also.

What is TREXIMET?

TREXIMET is a prescription medicine that contains sumatriptan and naproxen sodium (an NSAID). TREXIMET is used to treat acute migraine headaches with or without aura in patients 12 years of age and older.

TREXIMET is not used to treat other types of headaches such as hemiplegic (that make you unable to move on one side of your body) or basilar (rare form of migraine with aura) migraines.

TREXIMET is not used to prevent or decrease the number of migraine headaches you have.

It is not known if TREXIMET is safe and effective to treat cluster headaches.

Who should not take TREXIMET?

Do not take TREXIMET if you have:

- heart problems, history of heart problems, or right before or after heart bypass surgery
- had a stroke, transient ischemic attack (TIAs), or problems with your blood circulation
- hemiplegic migraines or basilar migraines. If you are not sure if you have these types of migraines, ask your healthcare provider.
- narrowing of blood vessels to your legs and arms (peripheral vascular disease), stomach (ischemic bowel disease), or kidneys
- taken any medicines in the last 24 hours that are called 5-HT₁ agonists that are triptans or contain ergotamine. Ask your healthcare provider for a list of these medicines if you are not sure.
- taken an antidepressant medicine called a monoamine oxidase (MAO) inhibitor within the last 2 weeks. Ask your healthcare provider for a list if you are not sure.
- had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAID medicine
- an allergy to sumatriptan, naproxen, or any of the ingredients in TREXIMET. See "What are the ingredients in TREXIMET?" below for a complete list of ingredients.

What should I tell my healthcare provider before taking TREXIMET?

Before you take TREXIMET, tell your healthcare provider about all of your medical conditions, including if you:

- have high blood pressure
- have asthma
- have high cholesterol
- have diabetes
- have kidney problems
- have liver problems
- have had epilepsy or seizures
- are not using effective birth control
- are pregnant, think you might be pregnant, or are trying to become pregnant. **TREXIMET should not be used by pregnant women during the third trimester of their pregnancy.**
- are breastfeeding or plan to breastfeed. The components of TREXIMET pass into your breast milk and may harm your baby. Talk with your healthcare provider about the best way to feed your baby if you take TREXIMET.
- liver problems
- third trimester of pregnancy
- uncontrolled high blood pressure

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

TREXIMET and certain other medicines can affect each other, causing serious side effects.

What should I avoid while taking TREXIMET?

TREXIMET can cause dizziness, weakness, or drowsiness. If you have these symptoms, do not drive a car, use machinery, or do anything where you need to be alert.

What are the possible side effects of TREXIMET?

TREXIMET may cause serious side effects. See "What is the most important information I should know about TREXIMET?" These serious side effects include:

- changes in color or sensation in your fingers and toes (Raynaud's syndrome)
- new or worse high blood pressure
- heart failure from body swelling (fluid retention)
- kidney problems including kidney failure
- low red blood cells (anemia)
- liver problems including liver failure
- asthma attacks in people who have asthma
- stomach and intestinal problems (gastrointestinal and colonic ischemic events). Symptoms of gastrointestinal and colonic ischemic events include:
 - sudden or severe stomach pain
 - weight loss
 - constipation or diarrhea
 - fever
 - stomach pain after meals
 - nausea or vomiting
 - bloody diarrhea
- problems with blood circulation to your legs and feet (peripheral vascular ischemia). Symptoms of peripheral vascular ischemia include:
 - cramping and pain in your legs or hips
 - feeling of heaviness or tightness in your leg muscles
 - burning or aching pain in your feet or toes while resting
 - numbness, tingling, or weakness in your legs
 - cold feeling or color changes in 1 or both legs or feet
- medication overuse headaches. Some people who use too many TREXIMET tablets may have worse headaches (medication overuse headache). If your headaches get worse, your healthcare provider may decide to stop your treatment with TREXIMET.
- serotonin syndrome. Serotonin syndrome is a rare but serious problem that can happen in people using TREXIMET, especially if TREXIMET is used with antidepressant medicines called SSRIs or SNRIs.
- Stop taking TREXIMET and call your healthcare provider right away if you have any of the following symptoms of serotonin syndrome:
 - changes in blood pressure
 - tight muscles
 - mental changes such as seeing things that are not there (hallucinations), agitation, or coma
 - fast heartbeat
 - high body temperature
 - trouble walking
- seizures. Seizures have happened in people taking sumatriptan, one of the ingredients in TREXIMET, who have never had seizures before. Talk with your healthcare provider about your chance of having seizures while you take TREXIMET.

The most common side effects of TREXIMET include:

- dizziness
- pain, discomfort, or stiffness in your neck, throat, jaw, or chest
- tingling or numbness in your fingers or toes
- dry mouth
- heartbeat problems
- feeling weak, drowsy, or tired
- nausea
- heartburn
- feeling hot
- muscle tightness

Stop TREXIMET and call your healthcare provider right away if you have any of the following symptoms:

- nausea that seems out of proportion to your migraine
- vomit blood
- yellow skin or eyes
- more tired or weaker than usual
- itching
- swelling of the arms, legs, hands, and feet
- sudden or severe stomach pain
- blood in your bowel movement or it is black and sticky like tar
- unusual weight gain
- flu-like symptoms
- diarrhea
- tenderness in your upper right side

These are not all of the possible side effects of TREXIMET. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

For more information, please see the complete Prescribing Information, including BOXED WARNINGS, and the Medication Guide at TREXIMET.com.